


Parity-Violating Electron Scattering and Q_{weak}

David S. Armstrong
College of William & Mary



Symmetry Tests in Nuclei and Atoms

Kavli Institute for Theoretical Physics, UCSB

Sept 19-23 2016



WILLIAM & MARY

CHARTERED 1693






Outline

- 1) Intro to Parity-Violating Electron Scattering (PVES)
- 2) Qweak:
 - first results on the proton's weak charge
 - prospects for final result
 - Sensitivity to new physics
- 3) Further Standard Model Tests with PVES:
Plans at JLab-12 GeV

A brief history of parity violation

1930s – weak interaction needed to explain nuclear β decay


1950s – parity violation in weak interaction;
V-A theory to describe ^{60}Co decay



1970s – neutral weak current observed
at Gargamelle

late 1970s – parity violation observed in electron scattering - SLAC E122

Parity-violating electron scattering



Proposed by Ya. B. Zeldovich JETP 36 (1959)

$$A_{PV} = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-} \propto \frac{\left| \begin{array}{c} \text{Diagram 1} \\ \gamma \\ \text{Diagram 2} \\ Z^0 \end{array} \right|}{\left| \begin{array}{c} \text{Diagram 3} \\ \gamma \end{array} \right|^2} \propto \frac{|M_Z|}{|M_\gamma|}$$

Electroweak interference

$$A_{PV} \propto \frac{G_F Q^2}{4\pi\alpha\sqrt{2}} \left(g_A^e g_V^T + \beta g_V^e g_A^T \right) \sim 10^{-4} Q^2 [\text{GeV}^2]$$

PARITY NON-CONSERVATION IN INELASTIC ELECTRON SCATTERING[☆]

C.Y. PRESCOTT, W.B. ATWOOD, R.L.A. COTTRELL, H. DeSTAEBLER, Edward L. GARWIN,
A. GONIDEC¹, R.H. MILLER, L.S. ROCHESTER, T. SATO², D.J. SHERDEN, C.K. SINCLAIR,
S. STEIN and R.E. TAYLOR

Stanford Linear Accelerator Center, Stanford University, Stanford, CA 94305, USA

J.E. CLENDENIN, V.W. HUGHES, N. SASAO³ and K.P. SCHÜLER

Yale University, New Haven, CT 06520, USA

M.G. BORGHINI

CERN, Geneva, Switzerland

Phys. Lett. 77B (1978)

K. LÜBELSMEYER

Technische Hochschule Aachen, Aachen, West Germany


and

W. JENTSCHKE

II. Institut für Experimentalphysik, Universität Hamburg, Hamburg, West Germany


Received 14 July 1978

We have measured parity violating asymmetries in the inelastic scattering of longitudinally polarized electrons from deuterium and hydrogen. For deuterium near $Q^2 = 1.6$ (GeV/c)² the asymmetry is $(-9.5 \times 10^{-5})Q^2$ with statistical and systematic uncertainties each about 10%.



Pioneering Experiment
SLAC E122

Deep-inelastic scattering from isoscalar target



Textbook Physics: High Energy Physics (D.H. Perkins)

SLAC E122 *cont'd*

Also critical test of parton model

Pivotal to establishing Weinberg-Salam-Glashow $SU(2) \times U(1)$ gauge theory

Techniques


Optically pumped electron source: rapid helicity reversal, integrate scattered flux

monitor & feedback to control electron beam fluctuations

Followed by:

1989: Mainz ^9Be
W. Heil et al.

1990: MIT/Bates ^{12}C
P.A. Souder et al.



SLAC Experiments

SLAC E122 – crucial confirmation of **WSG electroweak model**

- Electron-deuteron deep inelastic scattering
- High luminosity: photoemission from NEA GaAs cathode
- Rapid helicity-flip (sign of e- polarization)
- Polarimetry to determine beam polarization
- Magnetic spectrometer: backgrounds and kinematic separation

$$A_{PV} \sim 100 \pm 10 \text{ ppm}$$

$$\sin^2\theta_W = 0.20 \pm 0.03$$

SLAC E158 – 1999

- electron-electron scattering - purely leptonic interaction
- electron-electron weak attractive force had never before been measured!

$$A_{PV} \sim -131 \pm 14 \pm 10 \text{ ppb}$$


$$\sin^2\theta_W = 0.2403 \pm 0.0013$$

Weak Charges

Electroweak Lagrangian → Parity-Violating electron-quark term:

$$\mathcal{L}_{PV}^{EW} = \frac{G_F}{\sqrt{2}} \left[g_A^e (\bar{e} \gamma_\mu \gamma_5 e) \cdot \sum_q g_V^q (\bar{q} \gamma^\mu q) + g_V^e (\bar{e} \gamma_\mu e) \cdot \sum_q g_A^q (\bar{q} \gamma^\mu \gamma_5 q) \right]$$

$$C_{1q} = 2g_A^e g_V^q$$



-Electroweak Charges-

Particle	Electric Charge	Weak Vector Charge ($\sin^2 \theta_W \approx \frac{1}{4}$)
u	$+\frac{2}{3}$	$-2C_{1u} = +1 - \frac{8}{3} \sin^2 \theta_W \approx +\frac{1}{3}$
d	$-\frac{1}{3}$	$-2C_{1d} = -1 + \frac{4}{3} \sin^2 \theta_W \approx -\frac{2}{3}$
p(uud)	+1	$Q_W^p = 1 - 4 \sin^2 \theta_W \approx 0$
n(udd)	0	$Q_W^n = -1$

← Proton's Weak Charge ("accidental" suppression: enhanced sensitivity to new physics)


Qweak: Proton's weak charge

For forward angle scattering
at low Q^2 :

A_{PV} accesses Q_W^p

$$A_{PV} \equiv \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R} \rightarrow \frac{G_F Q^2}{4\pi\alpha\sqrt{2}} Q_W^p$$

Use four-fermion contact interaction to parameterize the effective PV electron-quark couplings (mass scale and coupling)



New physics:

$$\begin{aligned} \sigma &\propto |M_\gamma + M_Z + M_{\text{new}}|^2 \\ &\sim |M_\gamma|^2 + 2M_\gamma M_Z^* + 2M_\gamma M_{\text{new}}^* \end{aligned}$$


Planned 4% measurement of proton's weak charge - probes TeV-scale new physics

$$\frac{\Lambda}{g} \sim \left(\sqrt{2} G_F \Delta Q_W^p \right)^{-\frac{1}{2}} \sim O(\text{TeV})$$

Erlar, Kurylov, and Ramsey-Musolf, PRD 68, 016006 2003

Qweak: Proton's weak charge


Examples of TeV scale new physics that Q_{weak} would be sensitive to are:



Q_{weak} is also sensitive to MeV-GeV scale mediators such as:


Dark Photon:

- Astrophysical motivation, observed in positron data
- Might be linked to muon g-2 anomaly




Dark Parity Violation: (Davoudiasl, Lee, Marciano, arXiv 1402.3620)

- New source of low energy PV via mass mixing between Z and Z_d with observable consequences
- Complementary to direct search for heavy dark photons



SUSY “phase space”



Extracting the weak charge

$$A_{PV} = -\frac{G_F Q^2}{4\pi\alpha\sqrt{2}} [Q_w^p + B(\theta, Q^2)Q^2]$$


Hadron structure enters here: electromagnetic and electroweak form factors...

Reduced asymmetry more convenient:

$$A_{red} = \frac{A_{PV}}{A_0} \quad A_0 = -\frac{G_F Q^2}{4\pi\alpha\sqrt{2}}$$

One must extrapolate to $Q^2 = 0$.

We measure A_{phys}^{PV}
at $Q^2 = 0.025 \text{ GeV}^2$.




The previous strange form factor program (experiments at MIT/Bates, JLab and MAMI) allow us to subtract our hadronic contribution

Electroweak Radiative Corrections

In the Standard Model, the weak charge is *defined* at $Q^2 = 0, E = 0$.

$$Q_W^p = [\rho_{NC} + \Delta_e][1 - 4 \sin^2 \hat{\theta}_W(0) + \Delta'_e] + \square_{WW} + \square_{ZZ} + \square_{\gamma Z}$$



Full expression for Q_W^p has energy dependent corrections – need precise calculations

The \square_{WW} and \square_{ZZ} are well determined from pQCD ($\propto \frac{1}{q^2 - M_{W(Z)}^2 + i\epsilon}$)

The $\square_{\gamma Z}$ isn't pQCD friendly due to the photon leg ($\propto \frac{1}{q^2 + i\epsilon}$)

Electroweak Radiative Corrections


In the Standard Model, the weak charge is *defined* at $Q^2 = 0, E = 0$.

$$Q_W^p = [\rho_{NC} + \Delta_e][1 - 4 \sin^2 \hat{\theta}_W(0) + \Delta'_e] + \square_{WW} + \square_{ZZ} + \square_{\gamma Z}$$

Uncertainty from these corrections on *current* results is irrelevant.

$\square_{\gamma Z}$ contribution to Q_W^p (Qweak kinematics)

Gorchtein & Horowitz <i>PRL 102, 091806 (2009)</i>	0.0026 ± 0.0026
Sibirtsev, Blunden & Melnitchouk, Thomas <i>PRD 82, 013011 (2010)</i>	$0.0047^{+0.0011}_{-0.0004}$
Rislow & Carlson <i>PRD 83, 13007 (2011)</i>	0.0057 ± 0.0009
Gorchtein, Horowitz & Ramsey-Muslof <i>PRC 84, 015502 (2011)</i>	0.0054 ± 0.0020
Hall, Blunden, Melnitchouk, Thomas & Young <i>arXiv:1304:7877 (2013)</i> (calculation constrained by PVDIS data)	0.0052 ± 0.00043




Calculations are primarily dispersion theory type
error estimates can be firmed up with data!

Qweak: inelastic asymmetry data taken at $W \sim 2.3 \text{ GeV}$, $Q^2 = 0.09 \text{ GeV}^2$


Meeting PVES Challenges

- 180 μ A beam current (JLab record)
- High power cryogenic target
- Rapid helicity reversal (960 Hz)
- Small scattering angle: toroidal magnet, large acceptance
- 6 GHz detected rates: data-taking in integrating mode
- Radiation hard detectors
- Low noise 18-bit ADCs
- Exquisite control of helicity-correlated beam parameters
- Four different kinds of helicity reversal:
 - Rapid (Pockels cell at source)
 - Slow (insertable $\lambda/2$ plate)
 - Ultra slow (Wien-reversal, $g-2$ spin flip)
- Two independent high-precision beam polarimeters
- High resolution Beam Current monitors
- Dedicated Tracking system for kinematics determination

The Q_{weak} Apparatus



The Q_{weak} Apparatus



First result

Q_{weak} ran from Fall 2010 – May 2012 (Hall C at JLab)

Four distinct running periods:

- Hardware checkout (Fall 2010-January 2011)
- Run 0 (Jan-Feb 2011)
- Run 1 (Feb – May 2011)
- Run 2 (Nov 2011 – May 2012)

We have completed and unblinded the analysis of “Run 0”
(about 1/25th of our total dataset).

D. Androic *et al.* Phys. Rev. Lett. 111 (2013)141803.

$$A_{PV}^p = -279 \pm 35(\text{stat}) \pm 29(\text{sys}) \text{ ppb} \quad \langle Q^2 \rangle = 0.0250 \pm 0.0006 \text{ GeV}^2$$

$$\langle E_{\text{beam}} \rangle = 1155 \text{ MeV} \quad \theta_{\text{eff}} = 7.90^\circ$$

Good agreement with Standard Model prediction


Reduced Asymmetry

in the forward-angle limit ($\theta=0$)


$$A_0 = -\frac{Q^2 G_F}{4\sqrt{2}\pi\alpha}$$

$$\overline{A_{LR}^p} = \frac{A_{LR}}{A_0} \xrightarrow{\theta \rightarrow 0} [Q_W^p - Q^2 B(Q^2)]$$


4% of total data



The C_{1q} & the neutron's weak charge



The C_{1q} & the neutron's weak charge




Combining this result with the most precise atomic parity violation experiment we can also extract, for the first time, the neutron's weak charge:

$$Q_W^n = -0.975 \pm 0.010$$

$$Q_W^n(SM) = -0.9890$$

Qweak Run 2 – Quality of Data

(statistics only - not corrected for beam polarization, AI target windows, ΔQ^2 , etc.)



Expect final result early next year; will be statistics-limited

PVDIS at 6 GeV

Repeat of SLAC E122
to 5x higher precision (JLab, Hall A)

Sensitive to: C_{2u} & C_{2d}
(axial quark – vector electron)

Two kinematic settings:


$$A_{\text{exp}} = [-91.1 \pm 3.1(\text{stat.}) \pm 3.0(\text{syst.})] \times 10^{-6}$$

$$A_{\text{SM}} = -87.7 \times 10^{-6}$$

$$A_{\text{exp}} = [-160.8 \pm 6.4(\text{stat.}) \pm 3.1(\text{syst.})] \times 10^{-6}$$

$$A_{\text{SM}} = (-158.9 \pm 1.0) \times 10^{-6}$$

D. Wang *et al.* Nature **506** (2014)67



Qweak and PVDIS combined

Electron & quark compositeness or contact interaction limits*:

$$\Lambda^\pm = v \left[\frac{8\sqrt{5}\pi}{|(2C_{2u} - C_{2d})_{Q^2=0}|^\pm} \right]^{1/2}$$


$$v = \sqrt{\sqrt{2}/(2G_F)} = 246.22 \text{ GeV}$$

>5.8 TeV & >4.6 TeV
(constructive & destructive int. with SM)

c.f. HERA (ZEUS & H1) limits
>3.2 & >3.8 TeV on $e_V q_A$ term

c.f. ATLAS, PRD **87** 015010(2013)
>9.5 TeV & >12.1 TeV
in left-left isoscalar model

(need to assume all other contact interactions are zero; PVES does not need this assumption)



*convention of Eichten, Lane & Peskin
PRL **50**, 811 (1983)

Future: PVES at JLab in 12 GeV era

MOLLER - precision Standard Model test by measuring weak charge of electron in PV electron-electron scattering (revisit SLAC E158)

SOLID - precision Standard Model test by measuring PV DIS on deuteron: improved access to quark weak axial couplings C_{2q}

Large kinematic coverage: disentangle CSV and higher-twist effects

Elsewhere: **P2** experiment at Mainz/MAMI (\rightarrow *Kurt Aulenbacher's talk*)
improve Q_{weak} by factor of 2-3 at lower Q^2

MOLLER at 12 GeV

Parity-violating electron-electron scattering:
weak charge of electron

Update SLAC E158

$$A_{PV} = 35 \text{ ppb}$$

$$\text{Luminosity: } 3 \times 10^{39} \text{ cm}^2/\text{s}$$

75 μA 80% polarized beam

$$\delta(A_{PV}) = 0.73 \text{ ppb}$$

$$\delta(Q_e^w) = \pm 2.1 \% \text{ (stat.)} \pm 1.1 \% \text{ (syst.)}$$


$$\mathcal{L}_{e_1 e_2} = \sum_{i,j=L,R} \frac{g_{ij}^2}{2\Lambda^2} \bar{e}_i \gamma_\mu e_i \bar{e}_j \gamma^\mu e_j \quad \Rightarrow \quad \frac{\Lambda}{\sqrt{|g_{RR}^2 - g_{LL}^2|}} = 7.5 \text{ TeV}$$

LEP2 (g_{LR} and sum) mass scale sensitivity: ~ 5.2 and 4.4 TeV

MOLLER: Lepton compositeness (strong coupling) – 47 TeV

Sensitivity to: Doubly-charged scalar, heavy Z' , SUSY, dark Z ...


MOLLER and weak mixing angle




Reminder: at tree-level

$$Q_W^e = (1 - 4 \sin^2 \theta_W)$$


Higgs discovery at LHC allows firm prediction of MOLLER asymmetry in Standard Model



MOLLER apparatus



SOLID – accessing the C_{2q} 's



$$A_{\text{iso}} = \frac{\sigma^l - \sigma^r}{\sigma^l + \sigma^r} = - \left(\frac{3G_F Q^2}{\pi\alpha 2\sqrt{2}} \right) \frac{2C_{1u} - C_{1d}(1 + R_s) + Y(2C_{2u} - C_{2d})R_v}{5 + R_s}$$

Cahn and Gilman, PRD **17** 1313 (1978) polarized electrons on deuterium

$$R_s(x) = \frac{2S(x)}{U(x) + D(x)} \xrightarrow{\text{Large } x} 0$$

$$R_v(x) = \frac{u_v(x) + d_v(x)}{U(x) + D(x)} \xrightarrow{\text{Large } x} 1$$

$$C_{1u} = -\frac{1}{2} + \frac{4}{3} \sin^2 \theta_W \approx -0.19$$

$$C_{1d} = \frac{1}{2} - \frac{2}{3} \sin^2 \theta_W \approx 0.35$$

$$C_{2u} = -\frac{1}{2} + 2 \sin^2 \theta_W \approx -0.04$$

$$C_{2d} = \frac{1}{2} - 2 \sin^2 \theta_W \approx 0.04$$


$$Y = \frac{1 - (1 - y)^2}{1 + (1 - y)^2 - y^2 \frac{R}{R+1}}$$

$$R(x, Q^2) = \sigma^l / \sigma^r \approx 0.2$$

$$x \equiv x_{\text{Bjorken}}$$

$$y \equiv 1 - E'/E$$

SOLID – accessing the C_{iq} 's




Blue bands represent expected data:
Qweak (left) and PVDIS-6GeV (right)

Green bands are proposed SOLID PVDIS



SOLID – Large Acceptance Device

- **Moderate running times**
 - Large Acceptance
 - High Luminosity on LH2 & LD2
- **Better than 1% errors for small bins**
- **Kinematics:**
 - Large Q^2 coverage
 - x-range 0.25-0.75
 - $W^2 > 4 \text{ GeV}^2$



- **Requirements:**
 - Solenoid contains low energy backgrounds (Møller, pions, etc)
 - Baffling to cut backgrounds
 - Trajectories measured after baffles
 - Fast tracking—GEM, particle ID, calorimetry, and pipeline electronics
 - Precision polarimetry (0.4%) Compton and atomic hydrogen Moller

PVES Experiment Summary



Physics sensitivity from contact interaction (LEP2 convention, $g^2 = 4\pi$)


	precision	$\Delta \sin^2 \bar{\theta}_W(0)$	Λ_{new} (expected)
APV Cs	0.58 %	0.0019	32.3 TeV
E158	14 %	0.0013	17.0 TeV
Qweak I	19 %	0.0030	17.0 TeV
Qweak final	4.5 %	0.0008	33 TeV
PVDIS	4.5 %	0.0050	7.6 TeV
SoLID	0.6 %	0.00057	22 TeV
MOLLER	2.3 %	0.00026	39 TeV
P2	2.0 %	0.00036	49 TeV
PVES ^{12}C	0.3 %	0.0007	49 TeV

Jens Erler

Summary

- **Qweak**: First measurement of proton's weak charge, consistent with Standard Model, 25x more data soon to be released
- **Qweak and PVDIS** at 6 GeV: constraints on new physics
- **MOLLER and SOLID**: major programs after JLab upgrade
two complementary Standard Model tests.

*Thanks to the organizers for the kind invitation!
And thanks to you who stayed for my talk rather than....*



The Qweak Collaboration

97 collaborators 23 grad students
10 post docs 23 institutions





Institutions:

- 1 University of Zagreb
- 2 College of William and Mary
- 3 A. I. Alikhanyan National Science Laboratory
- 4 Massachusetts Institute of Technology
- 5 Thomas Jefferson National Accelerator Facility
- 6 Ohio University
- 7 Christopher Newport University
- 8 University of Manitoba,
- 9 University of Virginia
- 10 TRIUMF
- 11 Hampton University
- 12 Mississippi State University
- 13 Virginia Polytechnic Institute & State Univ
- 14 Southern University at New Orleans
- 15 Idaho State University
- 16 Louisiana Tech University
- 17 University of Connecticut
- 18 University of Northern British Columbia
- 19 University of Winnipeg
- 20 George Washington University
- 21 University of New Hampshire
- 22 Hendrix College, Conway
- 23 University of Adelaide
- 24 Syracuse University



D. Androic,¹ D.S. Armstrong,² A. Asaturyan,³ T. Averett,² J. Balewski,⁴ K. Bartlett,² J. Beaufait,⁵ R.S. Beminiwattha,⁶ J. Benesch,⁵ F. Benmokhtar,⁷ J. Birchall,⁸ R.D. Carlini,^{5,2} G.D. Cates,⁹ J.C. Cornejo,² S. Covrig,⁵ M.M. Dalton,⁹ C.A. Davis,¹⁰ W. Deconinck,² J. Diefenbach,¹¹ J.F. Dowd,² J.A. Dunne,¹² D. Dutta,¹² W.S. Duvall,¹³ M. Elaasar,¹⁴ W.R. Falk,⁸ J.M. Finn,² T. Forest,^{15,16} D. Gaskell,⁵ M.T.W. Gericke,⁸ J. Grames,⁵ V.M. Gray,² K. Grimm,^{16,2} F. Guo,⁴ J.R. Hoskins,² K. Johnston,¹⁶ D. Jones,⁹ M. Jones,⁵ R. Jones,¹⁷ M. Kargiantoulakis,⁹ P.M. King,⁶ E. Korkmaz,¹⁸ S. Kowalski,⁴ J. Leacock,¹³ J. Leckey,² A.R. Lee,¹³ J.H. Lee,^{6,2} L. Lee,¹⁰ S. MacEwan,⁸ D. Mack,⁵ J.A. Magee,² R. Mahurin,⁸ J. Mammei,¹³ J.W. Martin,¹⁹ M.J. McHugh,²⁰ D. Meekins,⁵ J. Mei,⁵ R. Michaels,⁵ A. Micherdzinska,²⁰ A. Mkrтчhyan,³ H. Mkrтчhyan,³ N. Morgan,¹³ K.E. Myers,²⁰ A. Narayan,¹² L.Z. Ndukum,¹² V. Nelyubin,⁹ H. Nuhait,¹⁶ Nuruzzaman,^{11,12} W.T.H van Oers,^{10,8} A.K. Opper,²⁰ S.A. Page,⁸ J. Pan,⁸ K.D. Paschke,⁹ S.K. Phillips,²¹ M.L. Pitt,¹³ M. Poelker,⁵ J.F. Rajotte,⁴ W.D. Ramsay,^{10,8} J. Roche,⁶ B. Sawatzky,⁵ T. Seva,¹ M.H. Shabestari,¹² R. Silwal,⁹ N. Simicevic,¹⁶ G.R. Smith,⁵ P. Solvignon,⁵ D.T. Spayde,²² A. Subedi,¹² R. Subedi,²⁰ R. Suleiman,⁵ V. Tadevosyan,³ W.A. Tobias,⁹ V. Tvaskis,^{19,8} B. Waidyawansa,⁶ P. Wang,⁸ S.P. Wells,¹⁶ S.A. Wood,⁵ S. Yang,² R.D. Young,²³ P. Zang,²⁴ and S. Zhamkochyan³

MOLLER: if SUSY seen at LHC...



MSSM sensitivity if light superpartners and large $\tan \beta$

$$P_R = (-1)^{3(B-L)+2s}$$