

**Possibly useful relations:**

$T(K) = T(^{\circ}C) + 273.15$	$T(^{\circ}F) = \frac{9}{5}T(^{\circ}C) + 32^{\circ}$	$\Delta L = \alpha L \Delta T$
$\Delta V = \beta V \Delta T$	$PV = nRT$	$PV = NkT$
$n = N/N_A$	$v_{\text{rms}} = \sqrt{\frac{3kT}{m}}$	$\overline{KE} = \frac{1}{2}mv^2 = \frac{3}{2}kT$
$Q = mc\Delta T$	$Q = mL_V$	$Q = mL_f$
$Q = mL_s$	$\frac{Q}{t} = \frac{\kappa A(T_2 - T_1)}{d}$	$\frac{Q}{t} = \sigma\epsilon A(T_{\text{hot}}^4 - T_{\text{cold}}^4)$
$W = P\Delta V$	monoatomic: $\Delta U = \frac{3}{2}nR\Delta T$	diatomic: $\Delta U = \frac{5}{2}nR\Delta T$
	$\Delta U = Q - W$	
Isochoric: $\Delta V = 0$	Isothermal: $\Delta T = 0$	Isobaric: $\Delta P = 0$
Adiabatic: $\Delta Q = 0$	$\epsilon = \frac{W}{Q_H} = 1 - \frac{Q_C}{Q_H}$	$\epsilon_{\text{Carnot}} = 1 - \frac{T_C}{T_H}$
$\text{COP}_{\text{ref}} = \frac{Q_C}{W}$	$\text{COP}_{\text{hp}} = \frac{Q_H}{W}$	
$S = (\frac{Q}{T})_{\text{reversible}}$	$\Delta S_{\text{closed}} \geq 0$	$S = k \ln W$
$W = (\# \text{ microstates})/\text{macrostate}$		
$F = \frac{k q_1  q_2 }{r^2}$	$\vec{F} = q\vec{E}$	$E = \frac{kQ}{r^2}$
$\vec{E} = \vec{E}_1 + \vec{E}_2 + \vec{E}_3 + \dots$	$V = \frac{PE}{q}$	$V = Ed$
$V = \frac{kQ}{r}$	$Q = CV$	$C = \kappa \frac{\epsilon_0 A}{d}$
$E_{\text{cap}} = \frac{1}{2}QV = \frac{1}{2}CV^2 = \frac{Q^2}{2C}$		$C_p = C_1 + C_2 + C_3 + \dots$
$\frac{1}{C_s} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$	$I = \frac{\Delta Q}{\Delta t}$	$I = nqAv_d$
$V = IR$	$R = \rho \frac{l}{A}$	$\rho = \rho_0(1 + \alpha\Delta T)$
$P = IV$	$P = I^2R$	$P = V^2/R$
$V = V_0 \sin(2\pi ft)$	$I = I_0 \sin(2\pi ft)$	$P = I_0V_0 \sin^2(2\pi ft)$
$P_{\text{av}} = \frac{1}{2}I_0V_0 = I_{\text{rms}}V_{\text{rms}}$	$I_{\text{rms}} = I_o/\sqrt{2}$	$V_{\text{rms}} = V_o/\sqrt{2}$
$R_S = R_1 + R_2 + R_3 + \dots$	$\frac{1}{R_p} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$	$P = E/t$
$\tau = RC$	$V = V_0 \exp^{-t/\tau}$	$V = V_0(1 - \exp^{-t/\tau})$

$$\begin{aligned}
F &= |q|vB \sin \theta & r &= \frac{mv}{qB} & m &= \frac{qB^2r^2}{2V} \\
v &= E/B & V_{\text{Hall}} &= Blv & F &= ILB \sin \theta \\
\tau &= NIAB \sin \theta & B &= \frac{\mu_o I}{2\pi r} & B &= \frac{\mu_o I}{2R} \\
B &= \mu_0 nI & \frac{F}{l} &= \frac{\mu_o I_1 I_2}{2\pi r} & \vec{B} &= \vec{B}_1 + \vec{B}_2 + \vec{B}_3 + \dots \\
\phi &= BA \cos \theta & \epsilon &= -N \frac{\Delta \phi}{\Delta t} & \epsilon &= NBA \omega \sin \omega t \\
\epsilon &= Blv & \frac{I_s}{I_p} &= \frac{V_p}{V_s} = \frac{N_p}{N_s} & c &= 1/\sqrt{\epsilon_0 \mu_0} \\
c &= \lambda f & c &= \frac{E}{B} & n &= c/v \\
I &= P/A & I_{\text{ave}} &= \frac{1}{2}c\epsilon_0 E_0^2 = \frac{1}{2\mu_0}cB_0^2 = \frac{1}{2\mu_0}E_0 B_0 & & \\
I_0 &= 2I_{\text{ave}} & \theta_{inc} &= \theta_{ref} & n_1 \sin \theta_1 &= n_2 \sin \theta_2 \\
\theta_C &= \sin^{-1} \frac{n_2}{n_1} & P &= 1/f & m &= -\frac{d_i}{d_o} = \frac{h_i}{h_o} \\
\frac{1}{d_i} + \frac{1}{d_o} &= \frac{1}{f} & f &= \frac{R}{2} & &
\end{aligned}$$

$$\begin{aligned}
R &= 8.314 \text{ J}/(\text{mol} \cdot \text{K}) & N_A &= 6.02 \times 10^{23} \text{ mol}^{-1} & k &= 1.38 \times 10^{-23} \text{ J/K} \\
\sigma &= 5.69 \times 10^{-8} \text{ J}/(\text{s} \cdot \text{m}^2 \cdot \text{K}^4) & & & & \\
1 \text{ cal} &= 4.186 \text{ J} & 1 \text{ kcal} &= 10^3 \text{ cal} & 1 \text{ Liter} &= 10^{-3} \text{ m}^3 \\
1 \text{ atm} &= 10^5 \text{ Pa} & |q_e| &= 1.60 \times 10^{-19} \text{ C} & k &= 9.0 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2 \\
\epsilon_0 &= 8.85 \times 10^{-12} \text{ C}^2/(\text{N} \cdot \text{m}^2) & & & \mu_0 &= 4\pi \times 10^{-7} \text{ T} \cdot \text{m}/\text{A} \\
c &= 3.00 \times 10^8 \text{ m/s} & & & &
\end{aligned}$$