

Please write your name or initials on each piece of paper you turn in.

1. [35 points total] Suppose we have a neutral spin-0 particle “ $s$ ” that can interact with itself by a  $\phi^3$  interaction and can also interact with electrons, and the interaction Lagrangian is

$$\mathcal{L}_{int} = -\frac{1}{3!} g \phi^3 - f \bar{\psi} \psi \phi ,$$

where  $\psi$  is the Dirac field for the electron,  $\phi$  is the real scalar field for particle “ $s$ ”, and  $f$  and  $g$  are real constants. Let the mass of the scalar be  $M$  and neglect the mass of the electron.

Draw the Feynman diagram [5 points] for the process

$$e^+ + e^- \rightarrow s + s .$$

Assume  $g \gg f$  so that diagrams of  $\mathcal{O}(f^2)$  can be ignored compared to diagrams of  $\mathcal{O}(fg)$ . Also assume that the total center-of-mass energy is at least  $2M$ . Calculate [30 points] the center-of-mass differential cross section  $d\sigma/d\Omega$ .

2. [35 points total] Suppose photons were massless scalar (spin-0) particles, and that the interaction Lagrangian density was

$$\mathcal{L}_{int} = -ig \bar{\psi} \gamma_5 \psi \phi$$

where  $g$  is a real constant,  $\psi$  is the electron field, and  $\phi$  is a real scalar field, describing the scalar particles which we will also call  $\phi$ .

Find and simplify the matrix element for the analog of Compton scattering. (The analog of Compton scattering would be the process  $\phi(k_1) + e^-(p_1) \rightarrow \phi(k_2) + e^-(p_2)$ .)

Break down the problem as

- Draw the (two) lowest order non-trivial Feynman diagrams for this process.
- Write down the scattering amplitude  $\mathcal{M}$  corresponding to the Feynman diagrams.
- Simplify the scattering amplitude. I believe it can be written as something like

$$\mathcal{M} = (const.) \left( \frac{1}{\omega_1} \pm \frac{1}{\omega_2} \right) \bar{u}(p_2) \not{k}_1 u(p_1) \equiv A \times \bar{u}(p_2) \not{k}_1 u(p_1) ,$$

where  $\omega_1$  and  $\omega_2$  are the energies of the incoming and outgoing  $\phi$ 's in the target rest frame (lab). Find  $A$ . (You are not asked, for this problem and for today, to continue after finding  $\mathcal{M}$ .)

3. [10 points] From analyzing the interaction Lagrangian of question 1, what are the mass dimensions of  $f$  and  $g$  (in our units where  $\hbar = 1$  and  $c = 1$ )?
4. [10 points] For the interaction Lagrangian of question 2,

$$\mathcal{L}_{int} = -ig \bar{\psi} \gamma_5 \psi \phi$$

where  $\psi$  is the electron field and  $\phi$  is a real scalar field, show that  $g$  is real (or not) if  $\mathcal{L}$  is hermitian.

5. [10 points] Given a two particle to two particle process where the incoming momenta are  $p_1$  and  $p_2$  and the outgoing momenta are  $p_3$  and  $p_4$ , one has  $p_1 + p_2 = p_3 + p_4$ . Define

$$s = (p_1 + p_2)^2, \quad t = (p_1 - p_3)^2, \quad u = (p_1 - p_4)^2 .$$

Show that

$$s + t + u = 0$$

for the case that all four particles are massless.