

1. [25 points total] Let ψ be a Dirac field, which we call the electron field, let ϕ be a neutral scalar field, and let V^μ be a neutral vector field. Consider the following Lagrangians,

a) $\mathcal{L} = \mathcal{L}_0 + e \bar{\psi} \gamma_\mu \psi V^\mu,$

b) $\mathcal{L} = \mathcal{L}_0 + i f \bar{\psi} \gamma_5 \psi \phi,$

c) $\mathcal{L} = \mathcal{L}_0 + h \bar{\psi} \psi \phi^2,$

where e , f , and h are constants and \mathcal{L}_0 is the free Lagrangian. For each Lagrangian, draw all the lowest nontrivial order Feynman diagrams that contribute to electron-positron scattering. (I believe there should be two diagrams in each case.) Using the Feynman diagrams as guidance, successfully write down the electron-positron scattering amplitudes for each case.

2. [35 points total] Consider a hypothetical decay $\Lambda^0 \rightarrow p + \pi^-$. For this problem, Λ^0 is a polarized spin-1/2 particle of mass m_Λ and momentum P ; p is a massless (this is where the “hypothetical” comes in) spin-1/2 proton of momentum p ; and π^- is a massless spin-0 particle of momentum k . Let the Lagrangian density be

$$\mathcal{L} = \bar{\psi}_p (g_V \gamma_\mu - g_A \gamma_\mu \gamma_5) \psi_\Lambda \times i \partial^\mu \phi + \text{hermitian conjugate} ,$$

where g_V and g_A are real constants, and ϕ is the field for the π^- .

(a) Draw the Feynman diagram for the decay.

(b) Write down the amplitude \mathcal{M} for the decay.

(c) Work the amplitude into the form

$$\mathcal{M} \propto m_\Lambda \bar{u}(p) (g_V + g_A \gamma_5) u(P) ,$$

where the polarization arguments are tacit.


(d) Sum $|\mathcal{M}|^2$ over the appropriate polarizations.

Query: Did you have the minus sign in $\{\bar{u}(p) (g_V + g_A \gamma_5) u(P, S)\}^\dagger = \bar{u}(P, S) (g_V - g_A \gamma_5) u(p)$?

Further small note: The case of real Λ^0 , real p , and real π^- is not so far from the above with $g_V \approx g_A$.

3. [20 points total] Evaluate the integral $\int d\alpha d\beta e^{\alpha M \beta}$ for the case that α and β are independent Grassmann variables, and M is an ordinary number (or a 1×1 matrix).

4. [20 points total] (a) For the three lowest energy levels of hydrogen, indicate the possible E1 and M1 radiative decays ($nL \rightarrow n'L' + \gamma$), labeling each as E1 or M1 as the case may be. To keep the diagram simpler, only draw decays where n and n' differ by one unit. Neglect energy differences among states with the same n .



(b) What do you expect is the main decay of the 3S state? Why?

(c) Of all the states in the diagram, which, if any, is (are) metastable? Why?