

Physics 722, Spring 2021

Problem Set 1, due Thursday, Feb 11.

1. Born Approximation and the Yukawa Potential

a) Consider the theory of a spinor field $\psi(x)$ coupled to a scalar field $\phi(x)$, with Lagrangian

$$\mathcal{L} = \bar{\psi} (i\not{\partial} - M) \psi + \frac{1}{2} (\partial_\mu \phi)^2 - \frac{m^2}{2} \phi^2 - g \bar{\psi} \psi \phi. \quad (1)$$

Compute the scattering amplitude for the process $\psi + \psi \rightarrow \psi + \psi$ to lowest nonvanishing order in the coupling g . By comparing the nonrelativistic limit of this scattering amplitude with the Born approximation to scattering in nonrelativistic quantum mechanics, determine the potential $V(\mathbf{x})$ corresponding to the Yukawa interaction.

b) Is the interaction attractive or repulsive? Explain.

2. Renormalization of Spinor Fields

In class we argued that in any Lorentz-invariant theory with a parity symmetry, the vacuum-to-one-particle matrix element of a Dirac spinor field operator $\Psi(x)$ is the same as in the free theory up to an overall rescaling that depends on the interactions.

We demonstrated this explicitly for the case that the one-particle state is spin-up in the rest frame of the particle. Repeat the argument in the case that the one-particle state is spin-down.

3. Show that for a Lorentz-scalar field $\phi(x)$,

$$\langle 0 | \phi(0) | \mathbf{k} \rangle = \langle 0 | \phi(0) | \mathbf{0} \rangle,$$

where $|\mathbf{0}\rangle$ is a one-particle state with vanishing spatial momentum and $|0\rangle$ is the vacuum. Recall that we used this relation in the derivation that the physical mass defined by the mass-shell condition $\omega_{\mathbf{k}}^2 - \mathbf{k}^2 = m^2$ in the interacting theory is the same as the location of the pole in the renormalized scalar-field two-point function.

Hint: Think about the unitary operation that converts $\phi(x)$ to $\phi(\Lambda^{-1}x)$ for Lorentz transformation Λ . How does $\phi(0)$ transform under a Lorentz transformation?