

# System of linear algebraic equations

Eugeniy E. Mikhailov

The College of William & Mary



Lecture 21

Notes

---

---

---

---

---

---

---

---

Eugeniy Mikhailov (W&M)

Practical Computing

Lecture 21

1 / 7

## Mobile problem


Suppose someone provided us with 6 weights and 3 rods. We need to calculate the positions of suspension points.

If system in equilibrium torque must be zero at any pivot point

$$\begin{aligned} w_1 x_1 - (w_2 + w_3 + w_4 + w_5 + w_6) x_2 &= 0 \\ w_3 x_3 - (w_4 + w_5 + w_6) x_4 &= 0 \\ w_5 x_5 - w_6 x_6 &= 0 \end{aligned}$$

We need 3 more equation. Let's fix length of the rods

$$\begin{aligned} x_1 + x_2 &= L_{12} \\ x_3 + x_4 &= L_{34} \\ x_5 + x_6 &= L_{56} \end{aligned}$$



Notes

---

---

---

---

---

---

---

---

Eugeniy Mikhailov (W&M)

Practical Computing

Lecture 21

2 / 7

## Mobile problem continued

Let's define  $w_{26} = w_2 + w_3 + w_4 + w_5 + w_6$  and  $w_{46} = w_4 + w_5 + w_6$

$$\begin{aligned} w_1 x_1 - w_{26} x_2 &= 0 \\ w_3 x_3 - w_{46} x_4 &= 0 \\ w_5 x_5 - w_6 x_6 &= 0 \\ x_1 + x_2 &= L_{12} \\ x_3 + x_4 &= L_{34} \\ x_5 + x_6 &= L_{56} \end{aligned}$$

$$\sum_j A_{ij} x_j = B_i \rightarrow \mathbf{Ax} = \mathbf{B}$$

Matlab has a lot of built in functions to solve problem of this form

$$\begin{pmatrix} w_1 & -w_{26} & 0 & 0 & 0 & 0 \\ 0 & 0 & w_3 & -w_{46} & 0 & 0 \\ 0 & 0 & 0 & 0 & w_5 & -w_6 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ L_{12} \\ L_{34} \\ L_{56} \end{pmatrix}$$

Notes

---

---

---

---

---

---

---

---

Eugeniy Mikhailov (W&M)

Practical Computing

Lecture 21

3 / 7

## Inverse matrix method

$$\mathbf{Ax} = \mathbf{B}$$

$$\mathbf{A}^{-1} \mathbf{Ax} = \mathbf{A}^{-1} \mathbf{B}, \quad \det(\mathbf{A}) \neq 0$$

### Analytical solution

$$\mathbf{x} = \mathbf{A}^{-1} \mathbf{B}$$

### Matlab first way (not the fastest)

$$\mathbf{x} = \text{inv}(\mathbf{A}) * \mathbf{B}$$

### Matlab second way (recommended)

$$\mathbf{x} = \mathbf{A} \setminus \mathbf{B}$$

Notes

---

---

---

---

---

---

---

---

Eugeniy Mikhailov (W&M)

Practical Computing

Lecture 21

4 / 7

## Recall the mobile problem

If  $w_1 = 20$ ,  $w_2 = 5$ ,  $w_3 = 3$ ,  $w_4 = 7$ ,  $w_5 = 2$ ,  $w_6 = 3$ ,  $L_{12} = 2$ ,  $L_{34} = 1$ , and  $L_{56} = 3$ , then  $w_{26} = 20$  and  $w_{46} = 12$ .

$$\begin{pmatrix} 20 & -20 & 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & -12 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & -3 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 2 \\ 1 \\ 1 \\ 3 \end{pmatrix}$$

Notes

---

---

---

---

---

---

---

---

## Matlab mobile solution

```
A=[ ...
20, -20, 0, 0, 0, 0; ...
0, 0, 3, -12, 0, 0; ...
0, 0, 0, 0, 2, -3; ...
1, 1, 0, 0, 0, 0; ...
0, 0, 1, 1, 0, 0; ...
0, 0, 0, 0, 1, 1; ...
]
B= [ 0; 0; 0; 2; 1; 3 ]
% 1st method
x=inv(A)*B
% 2nd method
x=A\B
```

```
x =
1.0000
1.0000
0.8000
0.2000
1.8000
1.2000
```

```
Check
>> A*x-B
1.0e-15 *
0
0
0
0
0.2220
0
```

Notes

---

---

---

---

---

---

---

---

## When do and when not to do inverse matrix

Solutions based on Inverse matrix calculations involve extra (unnecessary for solution) steps and thus are slower

```
>> A=rand(4000);
>> B=rand(4000,1);
>> tic; x=inv(A)*B; toc
Elapsed time is 54.831124 seconds.
>> tic; x=A\B; toc
Elapsed time is 19.822778 seconds.
```

However it is handy to calculate inverse matrix in advance if you solve  $Ax = B$  for different  $B$  with the same  $A$ .

```
>> tic; Ainv=inv(A); toc
Elapsed time is 58.304244 seconds.
>> B1=rand(4000,1); tic; x1=Ainv*B1; toc
Elapsed time is 0.048547 seconds.
>> B2=rand(4000,1); tic; x2=Ainv*B2; toc
Elapsed time is 0.048315 seconds.
```

Notes

---

---

---

---

---

---

---

---

## When do and when not to do inverse matrix

Notes

---

---

---

---

---

---

---

---