

System of linear algebraic equations

Eugeniy E. Mikhailov

The College of William & Mary



Lecture 21

Mobile problem

Suppose someone provided us with 6 weights and 3 rods. We need to calculate the positions of suspension points.

If system in equilibrium torque must be zero at any pivot point

$$w_1 x_1 - (w_2 + w_3 + w_4 + w_5 + w_6) x_2 = 0$$

$$w_3 x_3 - (w_4 + w_5 + w_6) x_4 = 0$$


$$w_5 x_5 - w_6 x_6 = 0$$

We need 3 more equation. Let's fix length of the rods

$$x_1 + x_2 = L_{12}$$

$$x_3 + x_4 = L_{34}$$

$$x_5 + x_6 = L_{56}$$



Mobile problem continued

Let's define $w_{26} = w_2 + w_3 + w_4 + w_5 + w_6$ and $w_{46} = w_4 + w_5 + w_6$

$$w_1 x_1 - w_{26} x_2 = 0$$

$$w_3 x_3 - w_{46} x_4 = 0$$

$$w_5 x_5 - w_6 x_6 = 0$$

$$x_1 + x_2 = L_{12}$$

$$x_3 + x_4 = L_{34}$$

$$x_5 + x_6 = L_{56}$$

$$\sum_j A_{ij} x_j = B_i \rightarrow \mathbf{Ax} = \mathbf{B}$$

Matlab has a lot of built in functions to solve problem of this form

$$\begin{pmatrix} w_1 & -w_{26} & 0 & 0 & 0 & 0 \\ 0 & 0 & w_3 & -w_{46} & 0 & 0 \\ 0 & 0 & 0 & 0 & w_5 & -w_6 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ L_{12} \\ L_{34} \\ L_{56} \end{pmatrix}$$

Inverse matrix method

$$\mathbf{Ax} = \mathbf{B}$$

$$\mathbf{A}^{-1}\mathbf{Ax} = \mathbf{A}^{-1}\mathbf{B}, \quad \det(\mathbf{A}) \neq 0$$

Analytical solution

$$\mathbf{x} = \mathbf{A}^{-1}\mathbf{B}$$

Matlab first way (not the fastest)

$$\mathbf{x} = \text{inv}(\mathbf{A}) * \mathbf{B}$$

Matlab second way (recommended)

$$\mathbf{x} = \mathbf{A} \setminus \mathbf{B}$$

Recall the mobile problem

If $w_1 = 20$, $w_2 = 5$, $w_3 = 3$, $w_4 = 7$, $w_5 = 2$, $w_6 = 3$, $L_{12} = 2$, $L_{34} = 1$, and $L_{56} = 3$, then $w_{26} = 20$ and $w_{46} = 12$.

$$\begin{pmatrix} 20 & -20 & 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & -12 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & -3 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 2 \\ 1 \\ 3 \end{pmatrix}$$

Matlab mobile solution

```
A=[ ...  
20, -20, 0, 0, 0, 0; ...  
0, 0, 3, -12, 0, 0; ...  
0, 0, 0, 0, 2, -3; ...  
1, 1, 0, 0, 0, 0; ...  
0, 0, 1, 1, 0, 0; ...  
0, 0, 0, 0, 1, 1; ...  
]
```

```
B= [ 0; 0; 0; 2; 1; 3 ]
```

```
% 1st method
```

```
x=inv(A)*B
```

```
% 2nd method
```

```
x=A\B
```

```
x =  
1.0000  
1.0000  
0.8000  
0.2000  
1.8000  
1.2000
```

Check

```
>> A*x-B  
1.0e-15 *  
0  
0  
0  
0  
0.2220  
0
```

When do and when not to do inverse matrix

Solutions based on Inverse matrix calculations involve extra (unnecessary for solution) steps and thus are slower

```
>> A=rand(4000);  
>> B=rand(4000,1);  
>> tic; x=inv(A)*B; toc  
Elapsed time is 54.831124 seconds.  
>> tic; x=A\B; toc  
Elapsed time is 19.822778 seconds.
```

However it is handy to calculate inverse matrix in advance if you solve $\mathbf{Ax} = \mathbf{B}$ for different \mathbf{B} with the same \mathbf{A} .

```
>> tic; Ainv=inv(A); toc  
Elapsed time is 58.304244 seconds.  
>> B1=rand(4000,1); tic; x1=Ainv*B1; toc  
Elapsed time is 0.048547 seconds.  
>> B2=rand(4000,1); tic; x2=Ainv*B2; toc  
Elapsed time is 0.048315 seconds.
```