

Ordinary Differential equations

Eugeniy E. Mikhailov

The College of William & Mary



Lecture 19

Notes

ODE definitions

An ordinary equation of order n has the following form

$$y^{(n)} = f(x, y, y', y'', \dots, y^{(n-1)})$$

x independent variable

$y^{(i)} = \frac{\partial^i y}{\partial x^i}$, the i th derivative of $y(x)$

f the force term

Notes

First order ODE example

Example

the acceleration of a body is the first derivative of velocity with respect to the time and equals to the force divided by mass

$$a(t) = \frac{dv}{dt} = v'(t) = \frac{F}{m}$$

$t \rightarrow x$ independent variable

$v \rightarrow y$

$F/m \rightarrow f$

And we obtain the canonical form

$$y^{(1)} = f(x, y)$$

for the first order ODE

Notes

n th order ODE transformation to the system of first order ODEs

$$y^{(n)} = f(x, y, y', y'', \dots, y^{(n-1)})$$

We define the following variables

$$y_1 = y, y_2 = y', y_3 = y'', \dots, y_n = y^{(n-1)}$$

$$\begin{pmatrix} y_1' \\ y_2' \\ y_3' \\ \vdots \\ y_{n-1}' \\ y_n' \end{pmatrix} = \begin{pmatrix} f_1 \\ f_2 \\ f_3 \\ \vdots \\ f_{n-1} \\ f_n \end{pmatrix} = \begin{pmatrix} y_2 \\ y_3 \\ y_4 \\ \vdots \\ y_n \\ f(x, y_1, y_2, y_3, \dots, y_n) \end{pmatrix}$$

We can rewrite n th order ODE as a system of first order ODEs

$$\vec{y}' = \vec{f}(x, \vec{y})$$

Notes

Cauchy boundary conditions

$$\vec{y}' = \vec{f}(x, \vec{y})$$

This is the system of n equations and thus requires n constraints.

With Cauchy boundary conditions, we specify $\vec{y}(x_0) = \vec{y}_0$
i.e. initial conditions

$$\begin{pmatrix} y_1(x_0) \\ y_2(x_0) \\ y_3(x_0) \\ \vdots \\ y_n(x_0) \end{pmatrix} = \begin{pmatrix} y_{1_0} \\ y_{2_0} \\ y_{3_0} \\ \vdots \\ y_{n_0} \end{pmatrix} = \begin{pmatrix} y_0 \\ y'_0 \\ y''_0 \\ \vdots \\ y_0^{(n-1)} \end{pmatrix}$$

Notes

Problem example

If acceleration of the particle is given, then we find the position of the particle as a function of time by solving

$$x''(t) = a$$

First, we need to **convert it to the canonical form**: the system of the first order ODEs.

- $t \rightarrow x$ time as independent variable
- $x \rightarrow y \rightarrow y_1$ particle position
- $v \rightarrow y' \rightarrow y_2$ velocity
- $a \rightarrow f$ acceleration as a force term

so

$$x'' = a \rightarrow y'' = f \rightarrow \vec{y}' = \vec{f}(x, \vec{y}) \rightarrow \begin{pmatrix} y'_1 \\ y'_2 \end{pmatrix} = \begin{pmatrix} y_2 \\ f \end{pmatrix}$$

We also need to provide the initial conditions:

position $x_0 \rightarrow y_{1_0}$ and velocity $v_0 \rightarrow y_{2_0}$

Notes

Euler's method

Let's, for simplicity, consider a simple first order ODE (notice the lack of the vector notation)

$$y' = f(x, y)$$

There is an exact way to write the solution

$$y(x_f) = y(x_0) + \int_{x_0}^{x_f} f(x, y) dx$$

The problem is that $f(x, y)$ depends on y itself. However, on a small interval $[x, x + h]$, we can assume that $f(x, y)$ is constant.

Then, we can use the familiar box integration formula. In application to the ODE, this is called the Euler's method.

$$y(x + h) = y(x) + \int_x^{x+h} f(x, y(x)) dx \approx y(x) + f(x, y(x))h$$

Notes

Euler's method continued

$$y(x + h) = y(x) + f(x, y)h$$

All we need is to split our interval into a bunch of steps of the size h and leap frog from the first x_0 to the next one $x_0 + h$, then to the $x_0 + 2h$, and so on.

Now, we can make an easy transformation to the vector case (i.e. the n th order ODE)

$$\vec{y}(x + h) = \vec{y}(x) + \vec{f}(x, \vec{y})h$$

Similarly to the boxes integration method, which is inferior in comparison to more advance methods, for example, the trapezoidal and Simpson's, **the Euler's method is very imprecise for a given h** and there are better ways.

Notes

Stability issues for the numerical solution

Let's have a look at the first order ODE

$$y' = 3y - 4e^{-x}$$


It has the following analytical solution

$$y = Ce^{3x} + e^{-x}$$

If the initial condition $y(0) = 1$, then the solution is

$$y(x) = e^{-x}$$

The `ode_unstable_example.m` script compares the numerical and the analytical solutions



It is clear that the numerical solution diverges from the analytical solution. The problem is in the round off errors. From a computer point of view, $y(0) = 1 + \delta$. Thus, $C \neq 0$ and the numerical solution diverges.

Do not trust the numerical solutions (regardless of a method) without proper consideration!

Navigation icons: back, forward, search, etc.

Notes

Notes

Notes

Notes
