

Electron Scattering from an almost Free Neutron

K. Griffioen

Undergrad Seminar Phys309

15 March 2005

The Nucleon

- The neutron and proton are two charge states of the ‘same’ particle: the nucleon.
- p: uud, two up quarks and a down quark
- n: udd, two down quarks and an up quark
- The nucleon also contains gluons, the messenger particles for the strong force, and quark-antiquark pairs, u, ubar, d, dbar, s, sbar.

Structure Functions

- Jefferson Lab is a large electron microscope
- Proton radius is about 1 fm
- Resolution is about 0.1 fm
- We scatter from quarks inside the nucleon
- $F_2(x) = \sum_i e_i^2 x f_i(x)$ is the scattering probability
- Here x is the fraction of the proton momentum carried by the struck quark, e is the quark charge, and f is the probability distribution for observing a quark with momentum fraction x .

Neutron Data Are Important...

...but hard to get

- Free neutrons decay in 15 minutes.

- Radioactivity!



- Zero charge makes it difficult to create a dense target
Magnetic bottle: $10^3 - 10^4$ n/cm² [TU München]
Typical proton target: $4 \cdot 10^{23}$ p/cm² [10 cm LH]

=> Alternative Solution: Deuterons and Helium-3.

BUT: potentially large (and not completely known) nuclear corrections: Kinematic smearing, Binding and off-shell effects, “EMC-effect”, final state interactions, coherent processes, non-nucleonic components of the wave function...

What can we do?

To learn more about the structure of the neutron, we can try two approaches:

- Study modifications of the neutron structure for bound neutrons in detail, to single out the best theoretical description of binding effects.
- Use the best possible approximation for a free neutron target: a neutron that is “barely” off-shell.

In both cases: Lepton scattering off the deuteron with simultaneous detection of a “backwards going” proton:


$D(e, e' p_s) X$

Relativity


- $E^2 = p^2 + m^2$ (speed of light $c=1$)
- E is the total energy, p is the momentum, and m is the mass (energy in rest frame) of a particle.
- $Q^2 = p^2 - E^2$ for virtual photon exchanged
- $W^2 = E^2 - p^2$ for the struck nucleon (after absorbing the virtual photon).

“Spectator Tagging”

$$p_n = (M_D \Delta E_S, \Delta \vec{p}_S); \quad \Delta_n = 2 \Delta \Delta_S$$



$$p_S = (E_S, \vec{p}_S); \quad \Delta_S = \frac{E_S \Delta \vec{p}_S \cdot \hat{q}}{M_D / 2}$$



$$W^2 = (p_n + q)^2 = p_n \cdot p_n + 2((M_D \Delta E_S) \Delta \vec{p}_n \cdot \vec{q}) \Delta Q^2$$

$$\Delta M^{*2} + 2M \Delta (2 \Delta \Delta_S) \Delta Q^2$$

$$x^* = \frac{Q^2}{2p_n \cdot q} \Delta \frac{Q^2}{2M \Delta (2 \Delta \Delta_S)}$$

Modification of Bound Neutrons - the $D(e,e'p_s)$ Experiment

- Experiment 94-102 at Jefferson Lab
- Run period “E6” in Hall B (CLAS)
- 5.75 GeV / 7 nA Electrons on a 5 cm long LD_2 target $\Rightarrow L=10^{34}/\text{cm}^2\text{s}$
- 8 calendar weeks in spring of 2002; 4.5 billion triggers
- CLAS-Collaboration and 2 Ph.D. students:
Dr. Alexei Klimenko and Cornel Butuceanu



CEBAF Large Acceptance Spectrometer

Schematic Diagram







Photo of Hall B - CLAS has been opened up for service work

Experimental Details




A typical event




Acceptance for protons in the backward hemisphere

Results: Momentum Distribution



Inclusive Scattering off a “free” Neutron - the BoNuS* Experiment



- Experiment 03-012 at Jefferson Lab in Hall B (CLAS)
- 4 and 6 GeV / 200 nA electrons impinging on a 10 cm long D₂ gas target (7 atm) => $L = 0.4 \cdot 10^{34} / \text{cm}^2 \text{s}$
- PAC-approved for 2 calendar months of running (2005/6)
- Old Dominion Univ., Jefferson Lab, Hampton Univ., William & Mary, James Madison Univ., and the CLAS collaboration




Radial TPC Prototype

* BoNuS = **B**arely **o**ff-shell **N**ucleon **S**cattering

BoNuS - Experimental Setup



Target-detector system for slow protons




- Thin-walled gas target (7 atm., room temperature)
- **Radial Time Projection Chamber (RTPC) with Gaseous Electron Multipliers (GEMs)**
- 2 Tesla longitudinal magnetic field (to suppress Moller electrons and to measure momentum)
- 3-dimensional readout of position and energy loss (“pads”)


Detector Parameters

- Geometric Acceptance
 - Sensitive over 148 deg. In phi, 20cm in Z.
- Momentum Acceptance
 - Protons from ~ 70 MeV/c
- Proton Identification (*next slide*)
- Vertex Z resolution $< \sim 10$ mm
- Track Momentum Resolution
- Track E information from dE/dx - *studying*
- Rate & Timing Issues

Proton ID by dE/dx & Curvature




100 MeV/c pion




100 MeV/c proton


RTPC - Concept



Cross sectional view



Foil supports



Electric field distribution

Voltage distribution



target

He Gas

Drift Cathode

GEMs

Pad Electrodes
4x5 mm²





4 cm

6 cm


Drift

40 60 69 mm

RTPC - GEMs



300-500 V, Gain 100-200



RTPC - Data Acquisition

- Alice TPC electronics (CERN) with Altro Chip
- 16 channels, 10 bit ADC with up to 25 MHz data rate
- 3-dimensional track reconstruction (using drift time information and 2 -dimensional location of readout pads)









Figure 1.1. ALTR0 Processing Chain

pRTPC w/ Inverter/Driver Cards

Ribbons
To
Readout
System





RTPC - 1st Prototype



Tracks from a test run at Triangle Universities Nuclear Lab (TUNL) with 100 MeV/c protons


RTPC - 2nd Prototype




Scale 1:1, 1/8 of the final RTPC (1/2 length, 1/4 of 360°)

Tested at TUNL, planned engineering run with CLAS


Production Model: Exploded View




Production Model



BoNuS Target (11/04)



Acceptance for Protons




6 GeV electron beam, 20 “ideal” days -> registered “events”

Scattered electron within CLAS fiducial cuts, proton above 60 MeV/c und 90°


“VIPs”: $p(\text{proton}) < 100 \text{ MeV/c}$, $\angle_{pq} > 110^\circ$

Proton reconstructed by the RTPC


Proton Tracks at TUNL




dE/dx Analysis from TUNL



dE/dx Analysis from TUNL




dE/dx Analysis from TUNL




Software Needs Work, too!

- DAQ: Interface w/CLAS data structure
- Event visualization – OK and improving
- Track Fitting - Straight Tracks OK and improving

- Need to incorporate
 - Vdrift(R) (ongoing)
 - B-Field
 - Lorentz angle
- Need Simulation and Simulated Data





R_{HIT} vs. Time Bin
(V_{drift} varies with R)



Curved e⁻ Drift
N.B.: nonuniform B

Expected Data



Systematics ~5%



← d/u

↑
Duality

The Future - 11 GeV

$D(e,e'p_s)$

BoNuS



Conclusions

- Proton tagging with deuterium targets is a fantastic technique for understanding properties of the nucleon
- BONUS is the first almost free nucleon target, which will allow measurements free of uncertainties due to nuclear physics.
- New technical developments have made this possible.
- Data taking will occur in Fall 2005 at Jlab!